PASJ: Publ. Astron. Soc. Japan , 1–??,
© 2008. Astronomical Society of Japan.

arXiv:astro-ph/0312652v1 31 Dec 2003

Photometric Observations of an SU UMa-type Dwarf Nova VW Coronae Borealis during Outbursts

Daisaku NOGAMI
Hida Observatory, Kyoto University, Kamitakara, Gifu 506-1314
nogami@kwasan.kyoto-u.ac.jp
Makoto UEMURA, Ryoko ISHIOKA, Taichi KATO
Department of Astronomy, Faculty of Science, Kyoto University, Sakyo-ku, Kyoto 606-8502
Jochen PIETZ
Nollenweg 6, 65510 Idstein, Germany

(Received 2002 0; accepted 2002 0)

Abstract

We report the photometric observations of an SU UMa-type dwarf nova VW CrB during two superoutbursts in 2001 and 2003 and a normal outburst in 2003. Superhumps with a period of 0.07287(1) d were observed during the 2003 superoutburst. The change rate of the superhump period was positive. During the normal outburst, there are some hint of modulation up to a 0.2-mag amplitude. However, any periodicity was not found. The recurrence cycles of the normal outburst and the superoutburst, and the distance were estimated to be ≥ 50 d, $270 \sim 500$ d, and 690^{+230}_{-170} pc, respectively. These recurrence cycles are usual values for an SU UMa-type dwarf nova having this superhump period. The superhump period of VW CrB was the longest among those of the SU UMa stars with positive derivatives of the superhump period. The coverage of our observations was, however, not enough, and the variation of the $P_{\rm SH}$ change rate of VW CrB is still unknown. A superhump regrowth and a brightening were seen near the end of the plateau phase. Measuring the deviation of the start timings of the brightening and the superhump regrowth (>2 days in VW CrB) will be a key to reveal the mechanism of these phenomena.

Key words: accretion, accretion disks — stars: novae, cataclysmic variables — stars: dwarf novae — stars: individual (VW CrB)

1. Introduction

This variable star was discovered by Antipin (1996), and first designated as Var21 CrB. He found 9 outbursts from the Moscow collection plates obtained between JD 2441805 and 2444042, and suggested its dwarf nova nature showing two kinds (long and short) of outbursts. Based on this report, Kazarovets, Samus (1997) gave the permanent variable-star name of VW CrB in the 73rd Name List. Novák (1997) caught superhumps with a period of 0.0743(6) d during a long outburst in 1997 May, establishing that VW CrB is an SU UMa-type dwarf nova. A spectrum taken by Liu et al. (1999) during an outburst showed a blue continuum having broad Balmer absorption lines with a central emission core and He II absorption, which is a typical spectrum for an outbursting dwarf nova. Na D absorption was also marginally detected.

VW CrB is identified with USNO B1.0 1231-0276740 ($B1 = 20.35, R1 = 19.55$). The ROSAT sources, 1RXP J160003+3311.0 and 2RXP J160003.9+331115 are X-ray counterparts. No infrared counterpart was found in the 2 Micron All Sky Survey (2MASS) (Hoard et al. 2002).

2. Observation

We observed VW CrB during three outbursts in 2001 June-July, 2003 April, and 2003 July-August at Kyoto University, Ouda Station, and Hida observatory in Japan, and in Erfstadt in Germany. The observation log is summarized in table 1.

The Kyoto frames and Ouda frames were processed by the PSF photometry package developed by one of the authors (TK) after dark-subtraction and flat-fielding. All the frames obtained at Hida were reduced by the aperture photometry package in IRAF[1], after de-biasing and flat-fielding. The Erfstadt frames were dark-subtracted, flat-fielded, and analyzed with a PSF photometry package.

The magnitudes of VW CrB in the Erfstadt frames were measured differentially from the local comparison star USNO B1.0 1232-0272156, which is the star of ID 2 ($V = 14.044(2)$; $B - V = 0.634(3)$) in the Arne Henden's sequence[2]. When the long-term light curve of the 2003 July outburst was drawn (figure 1), we shifted the zero point of these differential magnitudes to fit them with the visual estimations reported to VSNET (Kato et al. 2003d). The differential magnitudes in the other frames were adjusted to be in accordance with the Erfstadt data.

The 1-sigma error for each differential magnitude varied

[1] IRAF is distributed by the National Optical Astronomy Observatories for Research in Astronomy, Inc. under cooperative agreement with the National Science Foundation.

[2] ⟨ftp://ftp.nofs.navy.mil/pub/outgoing/aah/sequence/sumner/vwcrb.seq⟩

Table 1. Log of observations.

Date			HJD-2400000 Start–End	Exposure Time (s)	Frame Number	filter	Instrument†
2001	July	2	52093.105–52093.202	30	223	no	A
		3	52094.082–52094.214	30	147	no	B
			52094.109–52094.215	30	221	no	A
2003	April	9	52739.148–52739.261	10	550	R_c	C
		10	52740.151–52740.220	10	309	R_c	C
		12	52742.126–52742.196	30	56	R_c	C
2003	July	26	52847.024–52847.240	30	483	no	A
		28	52849.358–52849.499	60	125	no	D
		29	52850.350–52850.474	60	100	no	D
		31	52851.955–52852.137	20	490	no	E
			52851.979–52852.109	30	306	no	A
			52852.350–52852.475	60	109	no	D
	August	1	52853.364–52853.425	60	22	no	D
		2	52854.194–52854.205	30	7	no	A
			52854.350–52854.533	60	145	no	D
		3	52855.349–52855.483	60	63	no	D
		4	52856.352–52856.465	60	46	no	D
		5	52857.346–52857.471	60	69	no	D
		6	52858.356–52858.477	60	81	no	D
		7	52859.355–52859.476	60	64	no	D
		9	52861.338–52861.492	60	95	no	D
		10	52862.354–52862.469	60	69	no	D
		11	52863.345–52863.471	60	83	no	D
		12	52864.347–52864.470	60	75	no	D
		14	52866.373–52866.380	60	2	no	D

A: 30cm Tel. + SBIG ST-7E (Kyoto), B: 25cm Tel. + SBIG ST-7E (Kyoto), C: 60cm Tel. + PixelVision (SITe SI004AB) (Ouda), D: 20cm Tel. + SBIG ST-6B (Erfstadt), E: 60cm Tel. + PixCellent S/T 00-3194 (SITe 003AB) (Hida)

from 0.01 mag to over 0.1 mag, depending on the target brightness and the sky condition.

3. Result

In this section, we describe the observational results in detail in the order of the 2003 July-August outburst, the 2001 June-July outburst, and the 2003 April outburst, since the 2003 July-August outburst was best covered and, therefore, most informative.

3.1. The 2003 July-August Outburst

The whole light curve of the 2003 July-August outburst is exhibited in figure 1. This outburst was found by Mike Simonsen at JD 2452844.639. Our observations were started about 2.4 days later than his report to VSNET, and within 4 days after onset of the outburst. The outburst duration was 20 days. The dashed line represents the trend of exponential decline of 0.097(4) mag d^{-1} which was calculated by linear regression of our data between JD 2452847 and 2452856. From JD 2452857, the decline suddenly stopped, which is a 'brightening' just before the following rapid decline. This phenomenon has been first systematically investigated by Kato et al. (2003c) recently.

Our first run showed that superhumps had already fully developed by HJD 2452847.0 within 4 days from the outburst onset, which confirmed the SU UMa nature of this star. A period analysis by the Phase Dispersion Minimization (PDM) method Stellingwerf (1978) was performed for the data between JD 2452847 and 2452866 after subtraction of the general decline trend including the final 'brightening'. Figure 2 gives the resultant theta diagram, which clearly shows the best estimated superhump period (P_{SH}) to be 0.07287(1) d. The error was calculated by the Lafler-Kinman methods (Fernie 1989).

The daily evolution of the superhump is depicted in figure 3. The superhump amplitude was 0.24 mag on JD 2452847, and gradually decayed to 0.10 mag on JD 2452854. After that, the amplitude again grown to 0.14 mag by JD 2452858, though the large error due to thin clouds smeared the superhump profile on JD 2452859. A clear second peak is seen around the phase = 0.4 on JD 2452858.

The superhump maximum timings measured by eye



Fig. 1. The long-term light curve of the 2003 July-August outburst. The abscissa is HJD - 2452000, and the ordinate is Visual (CCD) magnitude. The open circles and the lower arrows are the visual observations and the upper limits reported to VSNET. The filled squares are daily average magnitudes of our data. This outburst lasted about 20 days. The dashed line represents the exponential decay trend of 0.097(4) mag d^{-1} which was obtained using the daily averaged magnitudes before HJD 2452857. It should be noted that VW CrB rose from the exponential decay after 14th day of this outburst.



Fig. 2. Theta diagram of a PDM analysis to measure the superhump period from the data between JD 2452847 and 2452866. The abscissa is the frequency in the unit of cycle d^{-1}. The best estimated superhump period is 0.07287(1) d.

were listed in table 2. The cycle count (E) was set to be 1 at the first superhump maximum. Linear regression yields a following equation on the maximum timings:

$$C1 = 7.294(\pm0.005) \times 10^{-2} \cdot E - 0.019(\pm0.004). \quad (1)$$

The difference of the observed timings from the calculated ones systematically changed as shown in figure 4. A quadratic polynomial fitting to $O-C1$ yields:

$$\begin{aligned} C2 = {} & 3.39(\pm0.31) \times 10^{-6} \cdot (E-70)^2 \\ & +2.2(\pm1.7) \times 10^{-5} \cdot (E-70) \\ & -0.0077(\pm0.0008). \end{aligned} \quad (2)$$

The fittings were carried out using the non-linear least-squares Marquardt-Levenberg algorithm. The equation (2) indicates a positive derivative of the superhump maximum timings, meaning that the superhump period became longer. The change rate of the superhump period is $\dot{P}_{\rm SH}/P_{\rm SH} = 9.3(\pm0.9) \times 10^{-5}$. Note that these analyses are based on the data obtained by JD 2452857 and the large error during the 'brightening' phase unfortunately made the behavior of the superhump period left unclear.

3.2. The 2001 June-July Outburst

The long-term light curve of the 2001 July outburst is drawn in figure 5. The long duration of >15 d proves that this outburst was a superoutburst. Since the uncertainty of the zero points of each run is very large, ∼0.3 mag, it is difficult to judge whether the final 'brightening' occurred also in this superoutburst.

All the data points are plotted in figure 6. We could not detect any periodic modulations including superhumps, which is probably due to the large errors and the short



Fig. 3. Daily averaged superhump light curves folded by the superhump period of 0.07287 d. The superhumps had fully developed to the amplitude of 0.24 mag within 4 days from the onset of the superoutburst. The amplitude gradually became smaller to 0.10 mag till JD 2452854, then again developed to 0.14 mag on JD 2452858. The superhump profile was smeared by the large observation error owing to thin clouds on JD 2452859.

Table 2. Timings of the superhump maxima during the 2003 July-August superoutburst.

HJD−2452847	E	O−C1*
0.062(2)	1	0.0081
0.133(1)	2	0.0061
0.207(3)	3	0.0072
2.387(2)	33	−0.0010
2.456(1)	34	−0.0050
3.403(2)	47	−0.0062
5.008(2)	69	−0.0059
5.080(2)	70	−0.0068
5.371(2)	74	−0.0076
5.441(2)	75	−0.0105
7.424(4)	102	0.0031
9.392(3)	129	0.0017
10.425(3)	143	0.0136

* Using equation (1). The unit is day.



Fig. 4. $O-C$ diagram on the superhump maximum timings (see table 2 and equation (1)). The solid line is obtained by a quadratic polynomial fitting to $O-C1$ (equation (2)), indicating a positive derivative of the superhump period.



Fig. 5. Whole light curve of the 2001 June-July outburst. This superoutburst lasted at least 15 days. The zero point of our observations (open circles) have a large uncertainty of ∼0.3 mag. It is not clear whether the brightening just before the rapid decline occurred or not.



Fig. 6. Short-term light curve during the 2001 June-July outburst. The magnitudes are arbitrarily shifted each day. The filled squares and open circles represent the data obtained by a 30cm telescope and 25cm telescope, respectively. The large errors and the short coverages prevent us from judging existence of superhumps with an amplitude smaller than 0.13 mag.

coverages.

3.3. The 2003 April Outburst

Figure 7 displays the long-term light curve of the 2003 April outburst. Since the duration of this outburst was shorter than 5 days, it was clearly a normal outburst.

The light curve on JD 2452739 is shown in figure 8a. There are some hints of modulations. After subtracting the declining trend of 0.7 mag d^{-1}, we performed a PDM period analysis on the data. The resultant Θ diagram is figure 8b, where no strong signal of periodicity is seen. If we make a light curve folded by the superhump period after subtraction of the general decline trend, a plausible light curve emerges (figure 8c). This periodicity is, however, most doubtful, since the Θ diagram in figure 8b does not point to that frequency, the full amplitude of 0.04 mag is nearly equal to the 1-sigma error for each point, and the coverage of the data is not enough for the period.

4. Discussion

4.1. Outburst Behavior

The outbursts reported to VSNET are summarized in table 3.

VW CrB has been relatively well monitored. The maximum magnitude of the superoutburst is rather bright, reaching $m_{\rm vis} = 14.0$, and the superoutburst has long durations up to >23 d for an SU UMa-type dwarf nova.



Fig. 7. Whole light curve of the 2003 April outburst. The outburst duration was shorter than 5 days, suggesting a normal outburst.

For these reasons, all the superoutbursts has been likely caught after the start of monitoring by amateurs, except for during the season when this variable is not observable. Although the recurrence cycle of the superoutburst (supercycle, $T_{\rm s}$) is not stable, it is estimated to be $T_{\rm s} = 270 \sim 500$ d, assuming that one superoutburst was missed between the 1998 September superoutburst and the 2000 September one.

The normal outburst is $m_{\rm vis} = 14.4$ at the maximum, fainter than in the superoutburst, and the duration is shorter than 5 d. Then, some normal outbursts have been missed. Taking also into account the outbursts observed by Antipin (1996), it is proper to estimate the cycle of the normal outburst ($T_{\rm n}$) to be 50 d or more.

This supercycle and the ratio of the normal-outburst cycle to the supercycle are ordinary values for a dwarf nova with $P_{\rm SH} = 0.072$ d (see Warner 1995; Nogami et al. 1997b; Kato et al. 2003c).

The superoutburst duration, up to >23 d as mentioned before, is ≥1.5 times longer than those of usual SU UMa stars, and is comparable of those of ER UMa (Kato, Kunjaya 1995; Robertson et al. 1995) and the durations of the main outburst (until the dip) of superoutbursts in WZ Sge stars (e.g. the 2001 superoutburst of WZ Sge, see Ishioka et al. 2002; Patterson et al. 2002). In the current scheme of the standard thermal-tidal disk instability model (for a review, see Osaki 1996), the outburst enter the rapid decay phase when the surface density becomes smaller than the critical value at the outer edge of the contracting accretion disk and the cooling wave starts. There are three possible conditions to make the outburst duration longer in the disk instability model: 1) large mass is supplied during the outburst (as the explanation for the ER UMa case, Osaki 1995a), 2) larger mass is stored in the disk by the outburst ignition (as the explanation for the WZ Sge case, Osaki 1995b), and 3) the outburst lasts until the disk become smaller.

VW CrB, however, seems to have a normal mass trans-



Fig. 8. (a) Light curve taken on JD 2452739. (b) Theta diagram for the data shown in the upper panel after subtraction of the linear decline trend. Any strong periodicity was not suggested. The frequency of 13.7236 cycle d^{-1} is the superhump frequency. (c) Light curve folded by the superhump period after subtraction of the linear decline trend. Although a plausible modulation emerges, its reality is fairly weak.

Table 3. Previous outbursts.

Date*			JD	$V_{\rm max}$	D†	Type‡
1996	Jun.	26	50261	15.4		N
1996	Aug.	13	50309	14.3:	>6	S
1997	Feb.	27	50506	15.0:	4?	N
1997	May	11	50579	14.1	>20	S
1997	Aug.	20	50680	14.7	2	N
1998	Sep.	27	51083	13.8	>23	S
1999	Apr.	25	51293	14.4	1?	N
2000	Apr.	08	51643	14.6	>2	N
2000	Sep.	13	51800	13.9:	>11	S
2001	Jun.	18	52078	13.7	>15	S
2002	Jun.	23	52449	14.2	>18	S
2003	Apr.	09	52738	14.4	<5	N
2003	Jul.	24	52844	14.0	20	S

* The discovery date.
† Duration of the outburst in a unit of day.
‡ N: normal outburst, S: superoutburst.

fer rate, judging from its ordinary $T_{\rm s}$, and our data provides no hint of enhancement of the mass transfer rate during the superoutburst. The possibility 1) is hence rejected.

Concerning the possibility 2), the amount of the mass stored by the outburst ignition is related to physical parameters of the viscosity in quiescence, the disk radius, the primary mass (cf. Osaki 1995b). These parameters, however, also have an effect on $T_{\rm n}$ and $T_{\rm s}$. Since the cycle lengths are normal values for an SU UMa star, it is difficult to suppose unusual parameters on these parameters.

Therefore, the possibility 3) is most plausible. This possibility means that the tidal torque strongly works on the disk gas. Since the mass ratio dominantly determines the tidal effect, it is quite important to accurately measure the mass ratio of this system.

4.2. Distance

From the superhump period and the maximum magnitude of the normal outburst, we can estimate the distance to VW CrB in the same method as that used by Kato et al. (2002), Kato et al. (2003c) and Nogami et al. (2003a).

Warner (1987) proposed a relationship between the absolute magnitude at the outburst maximum and the orbital period $P_{\rm orb}$. The orbital period of VW CrB has not yet been measured, we substitute $P_{\rm SH}$ for $P_{\rm orb}$. There is a well-known, tight relationship between $P_{\rm orb}$ and $P_{\rm SH}$, and the error introduced by this substitution is negligible, compared with those due to other factors. Absence of the eclipse indicates that the inclination of this system is not so high, and the inclination effect to the observed flux (Warner 1986) should be very small. The absolute maximum magnitude is thus estimated to be $M_V = 5.20(\pm0.23)$, using the equation proposed by Warner (1987). This maximum magnitude should be compared by that of the normal outburst (Kato et al. 2002). Then, assuming the apparent maximum magnitude $m_V = 14.4(\pm0.4)$ from the data in table 3, the distance is guessed to be 690^{+230}_{-170} pc.

The proper motion of the star identified with the VW CrB is listed in the USNO B1.0 catalog as $(\mu_{\rm R.A.}, \mu_{\rm Dec}) = (-6(10), -26(7))$ in a unit of mas yr^{-1}. This distance is smaller than the secure upper limit estimated using this proper motion and the maximum expected velocity dispersion of CVs (Harrison et al. 2000).

The X-ray luminosity in the range of 0.5-2.5 keV of VW CrB is calculated to be $\log L_{\rm X} = 31.0 \pm 0.2$, using the ROSAT data of the X-ray counterparts and the distance in the same way by Verbunt et al. (1997). This luminosity is a little higher than, but not far from, the average value of SU UMa stars (see figures 6 and 8 in Verbunt et al. 1997).

4.3. Superhump Period Change

The superhump period had been believed to monotonically decrease, or stay constant at least (e.g. Warner 1985; Patterson et al. 1993), before the discovery of $P_{\rm SH}$ increase during the 1995 superoutburst in AL Com (Nogami et al. 1997a). Following this, some SU UMa stars have been confirmed to show similar behaviors: V485 Cen (Olech 1997), EG Cnc (Kato et al. 2003b), SW UMa (Semeniuk et al. 1997; Nogami et al. 1998), V1028 Cyg (Baba et al. 2000), WX Cet (Kato et al. 2001a), HV Vir (Kato et al. 2001b; Ishioka et al. 2003), V592 Cas (Kato, Starkey 2002), WZ Sge (Patterson et al. 2002; Ishioka et al. 2003, in preparation), EI Psc (Uemura et al. 2002), V1141 Aql (Olech 2003), and KS UMa (Olech et al. 2003). Superhump-period decrease was explained by shrinkage of the disk radius, and was regarded as a natural consequence of mass depletion from the disk (Osaki 1985). The mechanism of the $P_{\rm SH}$ increase phenomenon is, however, still an open question.

The superhump period of VW CrB is the longest among those of the SU UMa-type dwarf novae with positive $\dot{P}_{\rm SH}$, listed above. Noticing the concentration of these SU UMa stars near the orbital period minimum, Kato et al. (2001b) suggested that the period increase may be related to a low mass ratio and/or a low mass transfer rate. This may be supported by MN Dra (Nogami et al. 2003b). This star is considered to have a high mass ratio and a high mass transfer rate, based on its long $P_{\rm orb}$ of 0.105 d and short $T_{\rm s}$ of $\sim$60 d. And the $P_{\rm SH}$ derivative was $\dot{P}_{\rm SH}/P_{\rm SH} = 1.7 \times 10^{-3}$, about one order of magnitude larger than the largest value ever know.

Recent observations have revealed that some SU UMa stars with a relatively high mass ratio, a high mass transfer rate, and a long orbital period have $\dot{P}_{\rm SH}/P_{\rm SH} \sim 0$ (e.g. BF Ara, Kato et al. 2003a). Furthermore, Olech et al. (2003) most recently proposed that SU UMa stars show a positive $\dot{P}_{\rm SH}/P_{\rm SH}$ around the midst of the plateau phase, and then change $\dot{P}_{\rm SH}/P_{\rm SH}$ to be a minus value during a late phase of the superoutburst. Our present data at the late phase, unfortunately, contain a large error, and can not be useful to make clear the evolution of the superhump period throughout a superoutburst.

4.4. Final 'Brightening' and Superhump Regrowth

VW CrB gave rise to a phenomenon of superhump regrowth and a final brightening at the end of the plateau phase of the 2003 July-August superoutburst, as described in section 3.1. Kato et al. (2003c) did a literature survey, and discussed the relationship between the final brightening and the superhump regrowth. The former were seen in many short-$P_{\rm SH}$ systems (but few WZ Sge stars) and some long-$P_{\rm SH}$ systems with long $T_{\rm s}$. Dwarf novae having shown the latter feature, except for V725 Aql, are more concentrated to the short end of the $P_{\rm SH}$ distribution. Kato et al. (2003c) gave a remark that the appearance of superhump regrowth and brightening near the termination of the superoutburst are phenomenologically coupled.

VW CrB is located at a slightly longer side out of the $P_{\rm SH}$ region of dwarf novae with such phenomena, and has a relatively short $T_{\rm s}$ for a system with the brightening. This dwarf nova may be an important object to determine the critical value of the binary parameters separating the SU UMa stars with/without the final brightening and the superhump regrowth.

The superhump amplitude on JD 2452856 was already larger than on JD 2452854 (figure 3), while the brightening seemed to start on JD 2452856 (figure 1). To reveal the connection and the mechanism of the final brightening and the superhump regrowth, one of the keys will be more detailed observations in many SU UMa stars which measure deviation of the start timings of the brightening and the superhump regrowth.

5. Summary

The SU UMa nature of VW CrB was confirmed, and some parameters are determined: the superhump period of $P_{\rm SH}$ =0.07287(1) d, the $P_{\rm SH}$ change rate of $\dot{P}_{\rm SH}/P_{\rm SH} = 9.3(\pm0.9) \times 10^{-5}$, the supercycle of $T_{\rm s} = 270 \sim 500$ d, the cycle length of the normal outburst of $T_{\rm n} \geq 50$ d, the superoutburst duration up to >23 d, the distance of 690^{+230}_{-170} pc, and the X-ray (0.5-2.5 keV) luminosity of $\log L_{\rm X} = 31.0 \pm 0.2$.

The supercycle and the cycle length of the normal outburst are usual values for an SU UMa star. The superoutburst duration is very long, and comparable to that of ER UMa and the duration of the main outburst of superoutbursts in WZ Sge stars. This may be due to a strong tidal torque. It is important to gauge the mass ratio.

The superhump period of VW CrB was the longest among those of the SU UMa-type dwarf novae with positive derivatives of the superhump period. Although positive-$\dot{P}_{\rm SH}$ systems have been supposed to have short orbital periods, most recent observations show that the the sign of $P_{\rm SH}$ change rate varies even during one superoutburst in some SU UMa stars. Since our observations at the late phase of the 2003 superoutburst was performed under bad conditions, the variation of the $P_{\rm SH}$ change rate of VW CrB is a topic left for future observations.

During the 2003 superoutburst, we observed regrowth of the superhump amplitude and brightening near the end of the plateau phase. These features have been observed mainly in SU UMa stars having short superhump periods. The superhump period and the supercycle of VW CrB are slightly longer and shorter, respectively, than those in such systems. Measuring deviation of the start timings of the brightening and the superhump regrowth (>2 d in VW CrB) will help to reveal the mechanism of the superhump regrowth and the final brightening.

The authors are thankful to amateur observers for continuous reporting their valuable observations to VSNET. Thanks are also to the referee for the useful comments. This work is partly supported by a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists (MU and RI), and a grant-in-aid from the Japanese Ministry of Education, Culture, Sports, Science and Technology (No. 13640239, 15037205).

References

Antipin, S. V. 1996, Inf. Bull. Variable Stars, 4343
Baba, H., Kato, T., Nogami, D., Hirata, R., Matsumoto, K., & Sadakane, K. 2000, PASJ, 52, 429
Fernie, J. D. 1989, PASP, 101, 225
Harrison, T. E., McNamara, B. J., Szkody, P., & Gilliland, R. L. 2000, AJ, 120, 2649
Hoard, D. W., Wachter, S., Clark, L. L., & Bowers, T. P. 2002, ApJ, 565, 511
Ishioka, R. et al. 2003, PASJ, 55, 683
Ishioka, R. et al. 2002, A&A, 381, L41
Kato, T., Bolt, G., Nelson, P., Monard, B., Stubbings, R., Pearce, A., Yamaoka, H., & Richards, T. 2003a, MNRAS, 341, 901
Kato, T., & Kunjaya, C. 1995, PASJ, 47, 163
Kato, T., Matsumoto, K., Nogami, D., Morikawa, K., & Kiyota, S. 2001a, PASJ, 53, 893
Kato, T., Nogami, D., Matsumoto, K., & Baba, H. 2003b, PASJ, in press (astro-ph/0310426)
Kato, T., Nogami, D., Moilanen, M., & Yamaoka, H. 2003c, PASJ, in press (astro-ph/0307064)
Kato, T., Sekine, Y., & Hirata, R. 2001b, PASJ, 53, 1191
Kato, T., & Starkey, D. R. 2002, Inf. Bull. Variable Stars, 5358
Kato, T., Uemura, M., Ishioka, R., Nogami, D., Kunjaya, C., Baba, H., & Yamaoka, H. 2003d, PASJ, in press (astro-ph/0310209)
Kato, T., Uemura, M., Matsumoto, K., Kinnunen, T., Garradd, G., Masi, G., & Yamaoka, H. 2002, PASJ, 54, 999
Kazarovets, E. V., & Samus, N. N. 1997, Inf. Bull. Variable Stars, 4471
Liu, Wu., Hu, J. Y., Zhu, X. H., & Li, Z. Y. 1999, ApJS, 122, 243
Nogami, D., Baba, H., Kato, T., & Novák, R. 1998, PASJ, 50, 297
Nogami, D., Kato, T., Baba, H., Matsumoto, K., Arimoto, J., Tanabe, K., & Ishikawa, K. 1997a, ApJ, 490, 840
Nogami, D., Masuda, S., & Kato, T. 1997b, PASP, 109, 1114
Nogami, D. et al. 2003a, PASJ, in press (astro-ph/0310274)
Nogami, D. et al. 2003b, A&A, 404, 1067
Novák, R. 1997, Inf. Bull. Variable Stars, 4489
Olech, A. 1997, Acta Astron., 47, 281
Olech, A. 2003, Acta Astron., 53, 85
Olech, A., Schwarzenberg-Czerny, A., Kędzierski, P., Złoczewski, K., Mularczyk, K., & Wiśniewski, M. 2003, Acta Astron., 53, 175
Osaki, Y. 1985, A&A, 144, 369
Osaki, Y. 1995a, PASJ, 47, L11
Osaki, Y. 1995b, PASJ, 47, 47
Osaki, Y. 1996, PASP, 108, 39
Patterson, J., Bond, H. E., Grauer, A. D., Shafter, A. W., & Mattei, J. A. 1993, PASP, 105, 69
Patterson, J. et al. 2002, PASP, 114, 721
Robertson, J. W., Honeycutt, R. K., & Turner, G. W. 1995, PASP, 107, 443
Semeniuk, I., Olech, A., Kwast, T., & Nalezyty, M. 1997, Acta Astron., 47, 201
Stellingwerf, R. F. 1978, ApJ, 224, 953
Uemura, M. et al. 2002, PASJ, 54, 599
Verbunt, F., Bunk, W. H., Ritter, H., & Pfeffermann, E. 1997, A&A, 327, 602
Warner, B. 1985, in Interacting Binaries, ed. P. P. Eggelton & J. E. Pringle (Dordrecht: D. Reidel Publishing Company), 367
Warner, B. 1986, MNRAS, 222, 11
Warner, B. 1987, MNRAS, 227, 23
Warner, B. 1995, Ap&SS, 226, 187